

Bibliografia

Para todas as espécies

www.fao.org

www.iucnredlist.org

www.fishbase.org

www.sealifebase.org

- Cheung W.W.L., Pitcher T.J., Pauly D. (2005) A fuzzy logic expert system to estimate intrinsic extinction vulnerabilities of marine fishes to fishing. *Biological Conservation*, 124: 97–111.
- Fischer W., Bauchot M.-L., Schneider M. (1987a) Fiches FAO d'identification des espèces pour les besoins de la pêche (Révision 1). Méditerranée et mer Noire - Zone de pêche 37. Volume I: Végétaux et Invertébrés. Organisation des Nations Unies pour l'alimentation et l'agriculture (FAO), Rome, pp. 1–760.
- Fischer W., Bauchot M.-L., Schneider M. (1987b) Fiches FAO d'identification des espèces pour les besoins de la pêche (Révision 1). Méditerranée et mer Noire - Zone de pêche 37. Volume II: Vertébrés. Organisation des Nations Unies pour l'alimentation et l'agriculture (FAO), Rome, pp. 761–1530.
- Roper C.F.E., Sweeney M.J., Nauen C.E. (1984) FAO species catalogue. Vol. 3. Cephalopods of the world. An annotated and illustrated catalogue of species of interest to fisheries. FAO Fisheries Synopsis No. 125, 277pp.
- Santos M.N., Jiménez M.T., Abecasis D., Almeida C., Bentes L., Canas A., Coelho R., Ferreira M., Gonçalves J.S., Lino P.G., Muñoz J.L., Machado D., Monteiro P., Oliveira F., Pareja-Rios E., Ribeiro J., Robles V., Veiga P., Erzini K. (2007) Catálogo de especies de peces de interés comercial de la costa sur atlántica de la Península Ibérica. Santos M.N., Erzini K., Díaz A., Manzano C. (eds.). Instituto de Investigación y Formación Agraria y Pesquera. Viceconsejería, Andalucía, 145pp.

Choco-comum / Cuttlefish (*Sepia officinalis*)

- Gamito R., Costa M.J., Cabral, H. (2015) Fisheries in a warming ocean: trends in fish catches in the large marine ecosystems of the world. *Regional Environmental Change*, 15: 57–65.
- Gutowska M., Pörtner H.-O., Melzner F. (2008) Growth and calcification in the cephalopod *Sepia officinalis* under elevated seawater pCO₂. *Marine Ecology Progress Series*, 373: 303–309.

- Lacoue-Labarthe T., Martin S., Oberhänsli F., Teyssié J.-L., Jeffree R., Gattuso J.-P., Bustamante P. (2012) Temperature and pCO₂ affect the bioaccumulation of radionuclides and trace elements in the eggs of the common cuttlefish, *Sepia officinalis*. *Journal of Experimental Marine Biology and Ecology*, 413: 45–49.
- Pimentel M.S., Trübenbach K., Faleiro F., Boavida-Portugal J., Repolho T., Rosa, R. (2012) Impact of ocean warming on the early ontogeny of cephalopods: A metabolic approach. *Marine Biology*, 159: 2051–2059.
- Shaw P., Pérez-Losada M. (2000) Polymorphic Microsatellites in the Common Cuttlefish *Sepia Officinalis* (Cephalopoda). *Molecular Ecology*, 9: 237–8.

Polvo-comum / Octopus *(Octopus vulgaris)*

- Chédia J., Widien K., Amina B. (2010) Role of sea surface temperature and rainfall in determining the stock and fishery of the common octopus (*Octopus vulgaris*, Mollusca, Cephalopoda) in Tunisia. *Marine Ecology*, 31: 431–438.
- Gamito R., Pita C., Teixeira C., Costa M.J., Cabral H.N. (2016) Trends in landings and vulnerability to climate change in different fleet components in the Portuguese coast. *Fisheries Research*, 181: 93–101.
- Gamito R., Teixeira C.M., Costa M.J., Cabral, H.N. (2015) Are regional fisheries' catches changing with climate? *Fisheries Research*, 161: 207–216.
- Repolho T., Baptista M., Pimentel M.S., Dionísio G., Trübenbach K., Lopes V.M., Lopes A.R., Calado R., Diniz M., Rosa, R. (2014) Developmental and physiological challenges of octopus (*Octopus vulgaris*) early life stages under ocean warming. *Journal of Comparative Physiology B*, 184: 55–64.

Lula-comum / European squid *(Loligo vulgaris)*

- Coelho M.L., Quintela J., Bettencourt V., Olavo G., Villa H. (1994) Population structure, maturation patterns and fecundity of the squid *Loligo vulgaris* from Southern Portugal. *Fisheries Research*, 21: 87–102.
- Pecl G.T., Jackson G.D. (2008) The potential impacts of climate change on inshore squid: biology, ecology and fisheries. *Reviews in Fish Biology and Fisheries*, 18: 373–385.
- Rosa R., Trübenbach K., Pimentel M.S., Boavida-Portugal J., Faleiro F., Baptista M., Dionísio G., Calado R., Pörtner H.-O., Repolho T. (2014) Differential impacts of ocean acidification and warming on winter

and summer progeny of a coastal squid (*Loligo vulgaris*). Journal of Experimental Biology, 217: 518–525.

Pota-voadora / Southern shortfin squid *(Illex coindetii)*

Lefkaditou E., Politou C.-Y., Palialexis A., Dokos J., Cosmopoulos P., Valavanis V.D. (2008) Influences of environmental variability on the population structure and distribution patterns of the short-fin squid *Illex coindetii* (Cephalopoda: Ommastrephidae) in the Eastern Ionian Sea. In Developments in Hydrobiology 203: Essential Fish Habitat Mapping in the Mediterranean, pp. 71–90. Ed. by V.D. Valavanis. Springer, Dordrecht. 300pp.

González A.F., Castro B.G., Guerra A. (1996) Age and growth of the short-finned squid *Illex coindetii* in Galician waters (NW Spain) based on statolith analysis. ICES Journal of Marine Science, 53: 802–810.

González A.F., Guerra A. (1996) Reproductive biology of the short-finned squid *Illex coindetii* (cephalopoda, ommastrephidae) of the Northeastern Atlantic. Sarsia, 81: 107–118.

Vijai D. (2016) Egg masses of flying squids (Cephalopoda: Ommastrephidae). Journal of Shellfish Research, 35: 1007–1012.

Salman A. (2017) Fecundity and spawning strategy of shortfin squid *Illex coindetii* (Oegopsida: Ommastrephidae), in the Eastern Mediterranean. Turkish Journal of Fisheries and Aquatic Sciences, 7: 843–851.

Peixe-espada-branco / Silver scabbardfish *(Lepidopus caudatus)*

Dhaker T., Rimel B., Wafa H., Missaou, H. (2016) Reproductive biology of the silver scabbard fish *Lepidopus caudatus* (Euphrasen, 1788) in the East coast of Tunisia. Cahiers de Biologie Marine, 57: 95–101.

Tamboril-branco / Angler *(Lophius piscatorius)*

Afonso-Dias I.P., Hislop J.R.G. (1996) The reproduction of anglerfish *Lophius piscatorius* Linnaeus from the North-west coast of Scotland. *Journal of Fish Biology*, 49: 18–39.

Duarte R., Azevedo M., Landa J., Pereda P. (2001) Reproduction of anglerfish (*Lophius budegassa* Spinola and *Lophius piscatorius* Linnaeus) from the Atlantic Iberian coast. *Fisheries Research*, 51: 349–361.

Velasco F., Landa J., Barrado J., Blanco M. (2008) Distribution, abundance, and growth of anglerfish (*Lophius piscatorius*) on the Porcupine Bank (west of Ireland). *ICES Journal of Marine Science*, 65: 1316–1325.

Peixe-espada-preto / Black scabbardfish *(Aphanopus carbo)*

Afonso C., Lourenço H.M., Dias A., Nunes M.L., Castro M. (2007) Contaminant metals in black scabbard fish (*Aphanopus carbo*) caught off Madeira and the Azores. *Food Chemistry*, 101: 120–125.

Cruz C., Saraiva A., Santos M.J., Eiras J., Ventura C., Soares J.P., Hermida M. (2009) Parasitic infection levels by *Anisakis* spp. larvae (Nematoda: Anisakidae) in the black scabbardfish *Aphanopus carbo* (Osteichthyes: Trichiuridae) from Portuguese waters. *Scientia Marina*, 73: 115–120.

Neves A., Vieira A.R., Farias I., Figueiredo I., Sequeira V., Gordo L.S. (2019) Reproductive strategies in black scabbardfish (*Aphanopus carbo* Baixae, 1839) from the NE Atlantic. *Scientia Marina*, 73S2: 19–31.

Salmão-do-Atlântico / Atlantic salmon *(Salmo salar)*

Jonsson B., Jonsson N. (2004) Factors affecting marine production of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 61: 2369–2383.

Pescada-branca / European hake *(Merluccius merluccius)*

Hidalgo M., Massutí E., Moranta J., Cartes J., Lloret J., Oliver P., Morales-Nin B. (2008) Seasonal and short spatial patterns in European hake (*Merluccius merluccius* L.) recruitment process at the Balearic Islands (western Mediterranean): The role of environment on distribution and condition. *Journal of Marine Systems*, 71: 367–384.

- Hidalgo M., Tomás J., Høie H., Morales-Nin B., Ninnemann U.S. (2008) Environmental influences on the recruitment process inferred from otolith stable isotopes in *Merluccius merluccius* off the Balearic Islands. *Aquatic Biology*, 3: 195–207.
- Maravelias C.D., Tsitsika E.V., Papaconstantinou C. (2007) Environmental influences on the spatial distribution of European hake (*Merluccius merluccius*) and red mullet (*Mullus barbatus*) in the Mediterranean. *Ecological Research*, 22: 678–685.
- Sánchez F., Gil J. (2000) Hydrographic mesoscale structures and Poleward Current as a determinant of hake (*Merluccius merluccius*) recruitment in Southern Bay of Biscay. *ICES Journal of Marine Science*, 57: 152–170.

Congro / European conger *(Conger conger)*

- Correia A.T., Ramos A.A., Barros F., Silva G., Hamer P., Morais P., Cunha, R.L., Castilho R. (2012) Population structure and connectivity of the European conger eel (*Conger conger*) across the Norte-eastern Atlantic and western Mediterranean: integrating molecular and otolith elemental approaches. *Marine Biology*, 159: 1509–1525.

Robalo-legítimo / European seabass *(Dicentrarchus labrax)*

- Pope E.C., Ellis R.P., Scolamacchia M., Scolding J.W.S., Keay A., Chingombe P., Shields R.J., Wilcox R., Speirs D.C., Wilson R.W., Lewis C.N., Flynn K.J. (2014) European sea bass, *Dicentrarchus labrax*, in a changing ocean. *Biogeosciences*, 11: 2519–2530.
- Vinagre C., Ferreira T., Matos L., Costa M.J., Cabral H.N. (2009) Latitudinal gradients in growth and spawning of sea bass, *Dicentrarchus labrax*, and their relationship with temperature and photoperiod. *Estuarine, Coastal and Shelf Science*, 81: 375–380.
- Vinagre C., Madeira D., Narciso L., Cabral H.N., Diniz M.S. (2012) Impact of climate change on coastal versus estuarine nursery areas: cellular and whole-animal indicators in juvenile seabass *Dicentrarchus labrax*. *Marine Ecology Progress Series*, 464: 237–243.

Enguia-europeia / European eel *(Anguilla anguilla)*

- Durif C.M.F., Gjøsæter J., Vøllestad L.A. (2011) Influence of oceanic factors on *Anguilla anguilla* (L.) over the twentieth century in coastal habitats of the Skagerrak, Southern Norway. *Proceedings of the Royal Society of London B: Biological Sciences*, 278: 464–473.
- Henderson P.A., Plenty S.J., Newton L.C., Bird D.J. (2012) Evidence for a population collapse of European eel (*Anguilla anguilla*) in the Bristol Channel. *Journal of the Marine Biological Association of the United Kingdom*, 92: 843–851.
- ICES (2013) Report of the Joint EIFAAC/ICES Working Group on Eels (WGEEL), 18– 22 March 2013 in Sukarietta, Spain, 4–10 September 2013 in Copenhagen, Denmark. ICES CM 2013/ACOM: 18. 851pp.
- Kim J.-H., Rimbu N., Lorenz S.J., Lohmann G., Nam S.-I., Schouten S., Rühlemann C., Schneider R.R. (2004) North Pacific and North Atlantic sea-surface temperature variability during the Holocene. *Quaternary Science Reviews*, 23: 2141–2154.
- Miller M.J. (2009) Ecology of anguilliform leptocephali: remarkable transparent fish larvae of the ocean surface layer. *Aqua-BioScience Monographs*, 2: 1–94.
- Minegish Y., Aoyama J., Inoue J.G., Miya M., Nishida M., Tsukamoto K. (2005) Molecular phylogeny and evolution of the freshwater eels genus *Anguilla* based on the whole mitochondrial genome sequences. *Molecular Phylogenetics and Evolution*, 34: 134–46.
- Pacariz S., Westerberg H., Björk G. (2014) Climate change and passive transport of European eel larvae. *Ecology of Freshwater Fish*, 23: 86–94.

Corvinata-real / Squeteague *(Cynoscion regalis)*

- Lankford T.E., Targett T.E. (1994) Suitability of estuarine nursery zones for juvenile weakfish (*Cynoscion regalis*): effects of temperature and salinity on feeding, growth and survival. *Marine Biology*, 119: 611–620.

Pregado / Turbot *(Scophthalmus maximus)*

Imsland A.K., Foss A., Gunnarsson S., Berntssen M.H.G., FitzGerald R., Bonga S.W., Ham E., Nævdal G., Stefansson S.O. (2001) The interaction of temperature and salinity on growth and food conversion in juvenile turbot (*Scophthalmus maximus*). *Aquaculture*, 198: 353–367.

Nissling A., Johansson U., Jacobsson M. (2006) Effects of salinity and temperature conditions on the reproductive success of turbot (*Scophthalmus maximus*) in the Baltic Sea. *Fisheries Research*, 80: 230–238.

Sarrajão / Atlantic bonito *(Sarda sarda)*

Collette B.B. (2003) Family Scombridae Rafinesque 1815 – mackerels, tunas and bonitos. California Academy of Sciences Annotated Checklists of Fishes, No. 19, 28 p.

Macías D., Gómez-Vives M.J., García S., Ortiz de Urbina J.M. (2005) Reproductive characteristics of Atlantic bonito (*Sarda sarda*) from the South western Spanish Mediterranean. *Col. Vol. Sci. Pap. ICCAT*, 58: 470–483.

Sabatés A., Recasens L. (2001) Seasonal distribution and spawning of small tunas (*Auxis rochei* and *Sarda sarda*) in the Northwestern Mediterranean. *Scientia Marina*, 65: 95–100.

Valeiras J., Abad E. (2006) Capítulo 2.1.10.5: Bacoreta. In *Manual de ICCAT*, pp. 243–250.

Atum-rabilho / Atlantic bluefin tuna *(Thunnus thynnus)*

Muhling B.A., Lee S.-K., Lamkin J.T., Liu Y. (2011) Predicting the effects of climate change on bluefin tuna (*Thunnus thynnus*) spawning habitat in the Gulf of Mexico. *ICES Journal of Marine Science*, 68: 1051–1062.

Ravier C., Fromentin J.-M. (2004) Are the long-term fluctuations in Atlantic bluefin tuna (*Thunnus thynnus*) population related to environmental changes? *Fisheries Oceanography*, 13: 145–160.

Visbeck M.H., Hurrell J.W., Polvani L., Cullen H.M. (2001) The North Atlantic Oscillation: Past, present, and future. *Proceedings of the National Academy of Sciences of the United States of America*, 98: 12876–12877.

Raia-lenga / Thornback ray (*Raja clavata*)

Jones M.C., Dye S.R., Fernandes J.A., Frölicher T.L., Pinnegar J.K., Warren R., Cheung W.W. (2013) Predicting the impact of climate change on threatened species in UK waters. PLoS One, 8: e54216.

Carocho / Portuguese dogfish (*Centroscymnus coelolepis*)

Clarke M.W., Connolly P.L., Bracken, J.J. (2001) Aspects of reproduction of the deep water sharks *Centroscymnus coelolepis* and *Centrophorus squamosus* from west of Ireland and Scotland. Journal of the Marine Biological Association of the United Kingdom, 81: 1019–1029.

Veríssimo A., Gordo L., Figueiredo I. (2003) Reproductive biology and embryonic development of *Centroscymnus coelolepis* in Portuguese mainland waters. ICES Journal of Marine Science, 60: 1335–1341.

Veríssimo A., McDowell J.R., Graves J.E. (2011) Population structure of a deep-water squaloid shark, the Portuguese dogfish (*Centroscymnus coelolepis*). ICES Journal of Marine Science, 68: 555–563.

Tintureira / Blue shark (*Prionace glauca*)

Queiroz N., Lima F.P., Maia A., Ribeiro P.A., Correia J.P., Santos A.M. (2005) Movement of blue shark, *Prionace glauca*, in the north-east Atlantic based on mark-recapture data. Marine Biological Association of the United Kingdom, 85: 1107–1112.

Galhudo-malhado / Spiny dogfish (*Squalus acanthias*)

Taylor I.G., Gallucci V.F. (2009) Unconfounding the effects of climate and density dependence using 60 years of data on spiny dogfish (*Squalus acanthias*). Canadian Journal of Fisheries and Aquatic Sciences, 66: 351–366.

Pata-roxa / Lesser spotted dogfish *(Scyliorhinus canicula)*

Jeffree R.A., Warnau M., Teyssié J.-L., Markich S.J. (2006) Comparison of the bioaccumulation from seawater and depuration of heavy metals and radionuclides in the spotted dogfish *Scyliorhinus canicula* (Chondrichthys) and the turbot *Psetta maxima* (Actinopterygii: Teleostei). *Science of the Total Environment*, 368: 839–852.

Tubarão-albafar / Bluntnose sixgill shark *(Hexanchus griseus)*

Ebert D.A. (1986). Biological Aspects of the Sixgill Shark, *Hexanchus griseus*. *Copeia*, 1, 131-135.

<https://doi.org/10.2307/1444898>

Ebert D.A. (2002) Some observations on the reproductive biology of the sixgill shark *Hexanchus griseus* (Bonnaterre, 1788) from South African waters. *African Journal of Marine Science*, 24: 359–363.

Capap C, Hemida F, Lorget O, Barrull J, Mate I, Souissi JB, Bradae M.N. (2004) Reproductive biology of the Bluntnose sixgill shark *Hexanchus griseus* (Bonnaterre, 1788) (Chondrichthyes: Hexanchidae) from the Mediterranean Sea: a review. *ACTA ADRIAT*, 45 (1): 95-106,

Lavagante-europeu / Bluntnose sixgill shark *(Homarus gammarus)*

Arnold K.E., Findlay H.S., Spicer J.I., Daniels C.L., Boothroyd D. (2009) Effect of CO₂-related acidification on aspects of the larval development of the European lobster, *Homarus gammarus* (L.). *Biogeosciences*, 6: 1747–1754.

Schmalenbach I., Franke H.-D. (2010) Potential impact of climate warming on the recruitment of an economically and ecologically important species, the European lobster (*Homarus gammarus*) at Helgoland, North Sea. *Marine Biology*, 157: 1127–1135.

Small D.P., Calosi P., Boothroyd D., Widdicombe S., Spicer J.I. (2016) The sensitivity of the early benthic juvenile stage of the European lobster *Homarus gammarus* (L.) to elevated pCO₂ and temperature. *Marine Biology*, 163: 53.

Lagosta / Common spiny lobster *(Palinurus elephas)*

- Green B.S., Gardner C., Hochmuth J.D., Linnane A. (2014) Environmental effects on fished lobsters and crabs. *Reviews in Fish Biology and Fisheries*, 24: 613–638.
- Göni R, Daniel L. (2005) Review of the biology, ecology and fisheries of *Palinurus* spp. species of European waters: *Palinurus elephas* (Fabricius, 1787) and *Palinurus mauritanicus* (Gruvel, 1911). *Cahiers de Biologie Marine* (0007-9723) (Station Biologique de Roscoff), 2:127-142.
- Hunter 1999. Biology of The European Spiny Lobster, *Palinurus Elephas* (Fabricius, 1787) (Decapoda, Palinuridea). 72: 545–565. <https://doi.org/10.1163/156854099503609>

Lagostim / Norway lobster *(Nephrops norvegicus)*

- Sardà F. (1998) *Nephrops norvegicus* (L.): Comparative biology and fishery in the Mediterranean Sea. Introduction, conclusions and recommendations. *Scientia Marina* 62(S1): 5-15
- Chapter Two - Habitat and Ecology of *Nephrops norvegicus*
- Johnson M.P, Lordan C, Power A.M. 2013. Chapter Two - Habitat and Ecology of *Nephrops norvegicus*. *Advances in Marine Biology*. 64: 27-63.
- Tuck I.D, Chapman C. J., Atkinson R. J. A. 1997. Population biology of the Norway lobster, *Nephrops norvegicus* (L.) in the Firth of Clyde, Scotland – I: Growth and density, *ICES Journal of Marine Science*. 54: 125–135. <https://doi.org/10.1006/jmsc.1996.0179>

Navalheira / Velvet crab *(Necora puber)*

- Rastrick S.P., Calosi P., Calder-Potts R., Foggo A., Nightingale G., Widdicombe S., Spicer J.I. (2014) Living in warmer, more acidic oceans retards physiological recovery from tidal emersion in the velvet swimming crab, *Necora puber*. *Journal of Experimental Biology*, 217: 2499–2508.

Spicer J.I., Raffo A., Widdicombe S. (2007) Influence of CO₂-related seawater acidification on extracellular acid–base balance in the velvet swimming crab *Necora puber*. *Marine Biology*, 151: 1117–1125.

Santola / Spider crab *(Maja squinado)*

González-Gurriarán E., Freire J., Bernárdez C. (2002) Migratory patterns of female spider crabs *Maja Squinado* detected using electronic tags and telemetry. *Journal of Crustacean Biology*, 22: 91–97.

Hartnoll R.G. (1963) The biology of Manx spider crabs. *Proceedings of the Zoological Society of London*, 141: 423–496.

Verísimo P., Bernárdez C., González-Gurriarán E., Freire J., Muiño R., Fernández L. (2011) Changes between consecutive broods in the fecundity of the spider crab, *Maja brachydactyla*. *ICES Journal of Marine Science*, 68: 472–478.

Caranguejo-azul / Blue crab *(Callinectes sapidus)*

Vasconcelos P., Carvalho A.N., Piló D., Pereira F., Encarnação J., Gaspar M.B., Teodósio M.A. (2019) Recent and consecutive records of the Atlantic blue crab (*Callinectes sapidus* Rathbun, 1896): rapid westward expansion and confirmed establishment along the southern coast of Portugal. *Thalassas: An International Journal of Marine Sciences*, 35: 485–494.

Caranguejo-verde / Green crab *(Carcinus maenas)*

Bessa F., Baeta A., Martinho F., Marques S., Pardal M.A. (2010) Seasonal and temporal variations in population dynamics of the *Carcinus maenas* (L.): the effect of an extreme drought event in a southern European estuary. *Journal of the Marine Biological Association of the United Kingdom*, 90: 867–876.

Xaputa / Atlantic pomfret *(Brama brama)*

Zhang Z., Liu C., Ding H., Xie P., Ma X., Guo Y., Xie C. (2017) Reproductive biology of bream *Abramis brama* (L.) in the lower reaches of the Irtys River, China. *Chinese Journal of Oceanology and Limnology*, 35: 1471–1481.

Verdinho / Blue whiting *(Micromesistius poutassou)*

Coombs S.H., Hiby A.R. (1979) The development of the eggs and early larvae of blue whiting, *Micromesistius poutassou* and the effect of temperature on development. *Journal of Fish Biology*, 14: 111–123.

Hátún H., Payne M.R., Jacobsen J.A. (2009) The North Atlantic subpolar gyre regulates the spawning distribution of blue whiting (*Micromesistius poutassou*). *Canadian Journal of Fisheries and Aquatic Sciences*, 66: 759–770.

ICES (2013) Report of the ICES Advisory Committee 2012. ICES Advice, 2013. Book 9. 185pp.

Faneca / Pouting *(Trisopterus luscus)*

Tanner S.E., Fonseca V.F., Cabral H.N. (2009) Condition of 0-group and adult pouting, *Trisopterus luscus* L., along the Portuguese coast: evidence of habitat quality and latitudinal trends. *Journal of Applied Ichthyology*, 25: 387–393.

Carapau-negrão / Blue jack mackerel *(Trachurus picturatus)*

Garcia A., Pereira J., Canha Â., Reis D., Diogo H. (2015) Life history parameters of blue jack mackerel *Trachurus picturatus* (Teleostei: Carangidae) from north-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom*, 95: 401–410.

Vasconcelos J., Alves A., Gouveia E., Faria G. (2006) Age and growth of the blue jack mackerel, *Trachurus picturatus* Bowdich, 1825 (Pisces: Teleostei) off Madeira archipelago. *Life and Marine Sciences*, 23: 47–57.

Vasconcelos J., Faria G., Freitas R., Gordo L.S. (2017) Fecundity regulation strategy of the blue jack mackerel, *Trachurus picturatus* (Bowdich, 1825), off Madeira Island (NE Atlantic). *Fisheries Research*, 190: 150–156.

Bica / Common pandora *(Pagellus erythrinus)*

Giannakopoulou L., Neofitou C. (2014) Heavy metal concentrations in *Mullus barbatus* and *Pagellus erythrinus* in relation to body size, gender, and seasonality. *Environmental Science and Pollution Research*, 21: 7140–7153.

Klimogianni A., Koumoundouros G., Kaspiris P., Kentouri M. (2004) Effect of temperature on the egg and yolk-sac larval development of common pandora, *Pagellus erythrinus*. *Marine Biology*, 145: 1015–1022.

Choupa / Black seabream *(Spondyliosoma cantharus)*

Gonçalves J.M.S., Erzini K. (2000) The reproductive biology of *Spondyliosoma cantharus* (L.) from the SW Coast of Portugal. *Scientia Marina*, 64: 403–411.

Pita C., Gamito S., Erzini K. (2002) Feeding habits of the gilthead seabream (*Sparus aurata*) from the Ria Formosa (southern Portugal) as compared to the black seabream (*Spondyliosoma cantharus*) and the annular seabream (*Diplodus annularis*). *Applied Ichthyology*, 18: 81–86.

Wassef E.A., Eisawy A. (1985) Food and feeding habits of wild and reared gilthead bream *Sparus aurata* L. *Cybium*, 9: 233–242.

Sargo-safia / Common two-banded seabream *(Diplodus vulgaris)*

Gonçalves J.M.S., Erzini, K. (2000) The reproductive biology of the two-banded sea bream (*Diplodus vulgaris*) from the southwest coast of Portugal. *Journal of Applied Ichthyology*, 16: 110–116.

Mouine N., Ktari M.-H., Chakroun-Marzouk N. (2010) Age and growth of *Diplodus vulgaris* (Sparidae) in the Gulf of Tunis. *Cybium*, 34: 37–45.

Besugo / Axillary seabream (*Pagellus acarne*)

Recasens L., Sabatés A., Demestre M., Martín P., Sanchez P. (2007) Relationship between temperature and fisheries: the case of *Pagellus acarne* in the NW Mediterranean. *Rapport du Comité International de la Mer Méditerranée*, 38: 580–581.

Ferreira / Sand steenbras (*Lithognathus mormyrus*)

Alssalam E.F.A., Ali S.M., El Sayed El Mor M.E.S., Ali R.A.S., Al Fergani E.S. (2016) Reproductive biology of the striped seabream *Lithognathus mormyrus* (Linnaeus, 1758) from Al Haneah Fishing Site, Mediterranean Sea, Eastern Libya. *Journal of Life Sciences*, 10: 171–181.

Tainha-garrento / Golden grey mullet (*Chelon auratus*)

Bat L., Şahin F., Öztekin A. (2018) Toxic metal amounts in *Chelon auratus* (Risso, 1810): a potential risk for consumer's health. *Journal of Aquaculture and Marine Biology*, 7: 303–306.

Dourada / Gilthead seabream (*Sparus aurata*)

Vargas-Chacoff L., Arjona F.J., Polakof S., Martín del Río M.P., Soengas J.L., Mancera J.M. (2009) Interactive effects of environmental salinity and temperature on metabolic responses of gilthead sea bream

Sparus aurata. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 154: 417–424.

Salmonete-legítimo / Surmullet *(Mullus surmuletus)*

Amin A.M., Madkour F.F., El-Regal M.A., Moustafa A.A. (2016) Reproductive biology of *Mullus surmuletus* (Linnaeus, 1758) from the Egyptian Mediterranean Sea (Port Said). International Journal of Environmental Science and Engineering, 7: 1–10.

Sargo-legítimo / White seabream *(Diplodus sargus)*

Leitão F., Baptista V., Teodósio M.A., Hughes S.J., Vieira V., Chícharo L. (2016) The role of environmental and fisheries multi-controls in white seabream (*Diplodus sargus*) artisanal fisheries in Portuguese coast. Regional Environmental Change, 16: 163–176.

Vinagre C., Cabral H.N., Costa M.J. (2010) Relative importance of estuarine nurseries for species of the genus *Diplodus* (Sparidae) along the Portuguese coast. Estuarine, Coastal and Shelf Science, 86: 197–202.

Cabra-cabaço / Tub gurnard *(Chelidonichthys lucerna)*

McCarthy I.D., Marriott A.L. (2018) Age, growth and maturity of tub gurnard (*Chelidonichthys lucerna* Linnaeus 1758; Triglidae) in the inshore coastal waters of Northwest Wales, UK. Journal of Applied Ichthyology, 34: 581–589.

Vallisneri M., Stagioni M., Montanini S., Tommasini S. (2011) Body size, sexual maturity and diet in *Chelidonichthys lucerna* (Osteichthyes: Triglidae) from the Adriatic Sea, north eastern Mediterranean. Acta Adriatica, 52: 141–148.

Calafate-de-riscas / Shi drum
(*Umbrina cirrosa*)

Boga-do-mar / Bogue
(*Boops boops*)

Sbrana A., Valente T., Scacco U., Bianchi J., Silvestri C., Palazzo L., Andrea de Lucia G., Valerani C., Ardizzone G., Matiddi M. (2020) Spatial variability and influence of biological parameters on microplastic ingestion by *Boops boops* (L.) along the Italian coasts (Western Mediterranean Sea). *Environmental Pollution*, 23: 114429.

Peixe-aranha-maior / Greater weever
(*Trachinus draco*)

Ak O., Genç A. (2013) Growth and reproduction of the greater weever (*Trachinus draco* L., 1758) along the eastern coast of the Black Sea. *Black Sea/Mediterranean Environment*, 19: 95–110.

Bagge O. (2004) The biology of the greater weever (*Trachinus draco*) in the commercial fishery of the Kattegat. *ICES Journal of Marine Sciences*, 61: 933–943.

Ordines F., Farriols M., Leonart J., Guijarro B., Quetglas A., Massutí E. (2014). Biology and population dynamics of by-catch fish species of the bottom trawl fishery in the western Mediterranean. *Mediterranean Marine Science*, 15: 613–625.

Šantić M., Pallaoro A., Rađa B., Jardas I. (2016). Diet composition of greater weever, *Trachinus draco* (Linnaeus, 1758) captured in the eastern-central Adriatic Sea in relation to fish size, season and sampling area. *Journal of Applied Ichthyology*, 32: 675–681.

Cabra-de-bandeira / Longfin gurnard
(*Chelidonichthys obscurus*)

Rascasso-escorpião / Small red scorpionfish *(Scorpaena notata)*

Muñoz M., Saabat M., Vila S., Casadevall M. (2005) Annual reproductive cycle and fecundity of *Scorpaena notata* (Teleostei: Scorpaenidae). *Scientia Marina*, 69: 555–562.

Ordines F., Quetglas A., Massutí E., Moranta J. (2009) Habitat preferences and life history of the red scorpion fish, *Scorpaena notata*, in the Mediterranean. *Estuarine, Coastal and Shelf Science*, 85: 537–546.

Charroco / Lusitanian toadfish *(Halobatrachus didactylus)*

Amorim M.C.P., Simões J.M., Almada V.C., Fonseca P.J. (2011) Stereotypy and variation of the mating call in the Lusitanian toadfish, *Halobatrachus didactylus*. *Behavioral Ecology and Sociobiology*, 65: 707–716.

Palazón-Fernández J.L., Arias A.M., Sarasquete C. (2001) Aspects of the reproductive biology of the toadfish, *Halobatrachus didactylus* (Schneider, 1801) (Pisces: Batrachoididae). *Scientia Marina*, 65: 131–138.

Carapau-branco / Atlantic horse mackerel *(Trachurus trachurus)*

Lavín A., Moreno-Ventas X., Ortiz de Zárate V., Abaunza P., Cabanas J.M. (2007) Environmental variability in the North Atlantic and Iberian waters and its influence on horse mackerel (*Trachurus trachurus*) and albacore (*Thunnus alalunga*) dynamics. *ICES Journal of Marine Science*, 64: 425–438.

Leitão F. (2015) Time series analyses reveal environmental and fisheries controls on Atlantic horse mackerel (*Trachurus trachurus*) catch rates. *Continental Shelf Research*, 111: 342–352.

Sarda / Atlantic mackerel *(Scomber scombrus)*

Borja A., Uriarte A., Egana J. (2002) Environmental factors and recruitment of mackerel, *Scomber scombrus* L. 1758, along the north-east Atlantic coasts of Europe. Fisheries Oceanography, 11: 116–127.

Cavala / Atlantic chub mackerel *(Scomber colias)*

Martins M.M., Skagen D., Marques V., Zwolinski J., Silva A. (2013) Changes in the abundance and spatial distribution of the Atlantic chub mackerel (*Scomber colias*) in the pelagic ecosystem and fisheries off Portugal. Scientia Marina, 77: 551–563.

Techetach M., Ajana R., Saoud Y. (2019) Reproductive parameters of Atlantic chub mackerel *Scomber colias* in M'diq Bay, Morocco. Journal of the Marine Biological Association of the United Kingdom, 99: 957–962.

Peixe-agulha / Garfish *(Belone belone)*

Ilhan D.U, Akalın S, Taskavak E, Toğulga M. 2004. Some biological characteristics of the garfish (*Belone belone* L., 1761) in Izmir Bay, Aegean Sea. Journal of Applied Ichthyology 20(5):413 – 416. DOI: 10.1111/j.1439-0426.2004.00592.x

Dorman J.A. (2006) Investigations into the biology of the garfish, *Belone belone* (L.), in Swedish waters. Journal of Fish Biology 39(1):59 – 69. DOI: 10.1111/j.1095-8649.1991.tb04341.x

Berbigão / Common edible cockle *(Cerastoderma edule)*

Morgan, E., O' Riordan, R. M., and Culloty, S. C. (2013). Climate change impacts on potential recruitment in an ecosystem engineer. Ecology and Evolution, 3: 581–594.

Dabouineau L, Ponsero A. 2009. Synthesis on biology of Common European Cockle *Cerastoderma edule*. Laurent Dabouineau, Alain Ponsero. Synthesis on biology of Common European Cockle *Cerastoderma edule*. 2009. ffhal-00581394f

Amêijoa-boia / Grooved carpet shell (*Ruditapes decussatus*)

Chícharo L., Chícharo M.A. (2001) Effects of environmental conditions on planktonic abundances, benthic recruitment and growth rates of the bivalve mollusc *Ruditapes decussatus* in a Portuguese coastal lagoon. *Fisheries Research*, 53: 235–250.

Trigui-El Ménif N., Le Pennec M., Maamouri F. (1995) La reproduction de la palourde *Ruditapes decussatus* (mollusque, bivalve) sur les côtes tunisiennes. *Vie Marine*, 5: 35–42.

Amêijoa-branca / Solid surf clam (*Spisula solida*)

Baptista V., Leitão F. (2014) Commercial catch rates of the clam *Spisula solida* reflect local environmental coastal conditions. *Journal of Marine Systems*, 130: 79–89.

Longueirão-direito / Sword razor shell (*Ensis siliqua*)

Croos M.E, Riordan R.M, Culloty S.C. 2014. The reproductive biology of the exploited razor clam, *Ensis siliqua*, in the Irish Sea. *Fisheries Research*. 150: 11-17

Conquilha / Truncate donax (*Donax trunculus*)

Gaspar M.B., Ferreira R., Monteiro C.C. (1999) Growth and reproductive cycle of *Donax trunculus* L., (Mollusca: Bivalvia) off Faro, southern Portugal. *Fisheries Research*, 41: 309–316.

Tirado C., Salas C. (1998) Reproduction and fecundity of *Donax trunculus* L., 1758 (Bivalvia: Donacidae) in the littoral of Malaga (Southern Spain). *Journal of Shellfish Research* 17: 169–176.

Ameijola / Smooth clam

(Callista chione)

Galimany E., Baeta M., Durfort M., Lleonart J., Ramón M. (2015) Reproduction and size at first maturity in a Mediterranean exploited *Callista chione* bivalve bed. *Scientia Marina*, 79: 233–242.

Amêijoa-japonesa / Japanese carpet shell *(Ruditapes philippinarum)*

Mexilhão / Mediterranean mussel *(Mytilus galloprovincialis)*

Lesser M.P. (2016) Climate change stressors cause metabolic depression in the blue mussel, *Mytilus edulis*, from the Gulf of Maine. *Limnology and Oceanography*, 61: 1705–1717.

Seuront L., Nicastro K.R., Zardi G.I., Goberville E. (2019) Decreased thermal tolerance under recurrent heat stress conditions explains summer mass mortality of the blue mussel *Mytilus edulis*. *Scientific reports*, 9: 1–14.

Zittier Z.M.C., Bock C., Lannig G., Pörtner H.-O. (2015) Impact of ocean acidification on thermal tolerance and acid–base regulation of *Mytilus edulis* (L.) from the North Sea. *Journal of Experimental Marine Biology and ecology*, 473: 16–25.

Gamba / Deep-water rose shrimp *(Parapenaeus longirostris)*

Ligas A., Sartor P., Colloca F. (2011) Trends in population dynamics and fishery of *Parapenaeus longirostris* and *Nephrops norvegicus* in the Tyrrhenian Sea (NW Mediterranean): the relative importance of fishery and environmental variables. *Marine Ecology*, 32: 25–35.

Camarão-vermelho / Blue and red shrimp *(Aristeus antennatus)*

Kapiris K., Thessalou-Legaki M. (2006) Comparative fecundity and oocyte size of *Aristaeomorpha foliacea* and *Aristeus antennatus* in the Greek Ionian Sea (E. Mediterranean) (Decapoda: Aristeidae). *Acta Zoologica*, 87: 239–245.

Kapiris K., Thessalou-Legaki M. (2009) Comparative reproduction aspects of the deep-water shrimps *Aristaeomorpha foliacea* and *Aristeus antennatus* (Decapoda, Aristeidae) in the Greek Ionian Sea (Eastern Mediterranean). *International Journal of Zoology*, Article ID 979512.

Camarão-branco-legítimo / Common prawn *(Palaemon serratus)*

Haig J., Ryan N.M., Williams K.F., Kaiser M.J. (2014) A review of the *Palaemon serratus* fishery: biology, ecology & management. Fisheries and Conservation Report, No. 38. Bangor University.

Silva T., Rangel M., Gonçalves M., Feijó D., Vian J., Felício M. (2006) Fecundity, embryonic development and mercury contamination of the *Palaemon serratus* at north coast of Portugal. Conference Abstract: IX Foro de los Recursos Maríños y de la Acuicultura de las Rías Gallegas. doi: 10.13140/RG.2.2.24646.52809

Trombeteiro / Longspine snipefish *(Macroramphosus scolopax)*

Marques V., Chaves C., Morais A., Cardador F., Stratoudakis Y. (2009) Distribution and abundance of snipefish (*Macroramphosus spp.*) off Portugal (1998-2003). *Scientia Marina*, 69: 563–576.

Pimpim / Boarfish *(Capros aper)*

White E., Minto C., Nolan C.P., King E., Mullins E., Clarke M. (2011) First estimates of age, growth, and maturity of boarfish (*Capros aper*): a species newly exploited in the Northeast Atlantic. *ICES Journal of Marine Science*, 68: 61–66.

Trombeiro / Blotched picarel (*Spicara maena*)

Dalouche F., Bensahla Talet L., Bensahla Talet A., Abi-Ayad S.M.E.A. (2019) Fecundity of the blotched picarel, *Spicara maena* (Linnaeus, 1758) from Oran Bay (Western Mediterranean Sea). *Journal of Fisheries*, 7(1): e309.

Labropoulou M., Papaconstantinou C. (2005) Effects of fishing on community structure of demersal fish assemblages. *Belgian Journal of Zoology*, 135: 191–197.

Sardinha / European pilchard (*Sardina pilchardus*)

Guisande C., Cabanas J.M., Vergara A.R., Riveiro I. (2001) Effect of climate on recruitment success of Atlantic Iberian sardine *Sardina pilchardus*. *Marine Ecology Progress Series*, 223: 243–250.

Santos M.B., González-Quirós R., Riveiro I., Cabanas J.M., Porteiro C., Pierce G.J. (2012) Cycles, trends, and residual variation in the Iberian sardine (*Sardina pilchardus*) recruitment series and their relationship with the environment. *ICES Journal of Marine Science*, 69: 739–750.

Biqueirão / European anchovy (*Engraulis encrasicolus*)

Bueno-Pardo J., Petitgas P., Kay S., Huret M. (2019) Integration of bioenergetics in an individual-based model to hindcast anchovy dynamics in the Bay of Biscay. *ICES Journal of Marine Science*, 77: 655–667.

Taboada F.G., Anadón R. (2016) Determining the causes behind the collapse of a small pelagic fishery using Bayesian population modeling. *Ecological Applications*, 26: 886–898.

Sardinela-da-Madeira / Madeiran sardinella (*Sardinella maderensis*)

Ba K., Thiaw M., Lazar N., Sarr A., Brochier T., Ndiaye I., Faye A., Sadio O., Panfili J., Thiaw T.O., Brehmer P. (2016) Resilience of key biological parameters of the Senegalese flat sardinella to overfishing and climate change. PLoS One, 11: e0156143.

Rodvalho / Brill
(*Scophthalmus rhombus*)

Linguado-legítimo / Common sole
(*Solea solea*)

El- Aiatt A.A.O, Shalloof K, El-Far A (2019). Reproductive Biology of the Common Sole, *Solea solea* in Southern East Mediterranean, Bardawil Lagoon, Egypt. Egyptian Journal of Aquatic Biology and Fisheries 23(1):403-411. DOI: 10.21608/ejabf.2019.29183

Azevia / Bastard sole
(*Microchirus azevia*)

Boufersaoui S., Handjar H. (2019) Population fecundity of seven fishes from Algerian coastal waters (South-Western Mediterranean Sea). Frontiers in Marine Sciences. Conference Abstract: XVI European Congress of Ichthyology. doi: 10.3389/conf.fmars.2019.07.00013

Percebe / Stalked barnacle
(*Pollicipes pollicipes*)

Cruz T., Araújo J. (1999) Reproductive patterns of *Pollicipes pollicipes* (Cirripedia: Scalpellomorpha) on the southwestern coast of Portugal. Journal of Crustacean Biology, 19: 260–267.